arXiv:1301.6531v1 [math.CO] 28 Jan 2013

JACK POLYNOMIALS AND THE MEASURE OF NON-ORIENTABILITY OF MAPS.
PART 1. ORIENTABILITY GENERATING SERIES

MACIEJ DOŁĘGA, VALENTIN FÉRAY, AND PIOTR ŚNIADY

ABSTRACT. We introduce the *orientability generating series* which measures non-orientability of a given map (i.e., a graph drawn on a surface). Presumably this quantity is the underlying structure behind the combinatorics of Jack polynomials and Jack characters and we formulate some concrete conjectures about this relationship. Our conjecture would give a positive answer to some positivity conjectures of Lassalle.

1. INTRODUCTION

1.1. **Jack polynomials and Macdonald polynomials.**

Henry Jack [Jac71] introduced a family of symmetric polynomials — which is now known as *Jack polynomials* $J_{\pi}^{(\alpha)}$ — indexed by an additional deformation parameter α. From the contemporary viewpoint probably the main motivation for studying Jack polynomials comes from the fact that they are a special case of the celebrated *Macdonald polynomials* which *“have found applications in special function theory, representation theory, algebraic geometry, group theory, statistics and quantum mechanics”* [GR05]. Indeed, some surprising features of Jack polynomials [Sta89] have led in the past to the discovery of Macdonald polynomials [Mac95] and Jack polynomials have been regarded as a relatively easy case [LV95] which later allowed understanding of the more difficult case of Macdonald polynomials [LV97]. A brief overview of the Macdonald polynomials (and their relationship to Jack polynomials) is given in [GR05].

In this article we will address some conjectures related to combinatorics of the (relatively simple) Jack polynomials in the hope that eventually this might lead to the better understanding of the general case of Macdonald polynomials.

2010 *Mathematics Subject Classification.* Primary 05E05; Secondary 05C10, 05C30, 20C30 .

Key words and phrases. Jack polynomials, Jack characters, maps, topological aspects of graph theory.



Figure 1.1. Example of an *oriented map*. The map is drawn on a torus: the left side of the square should be glued to the right side, as well as bottom to top, as indicated by arrows.

1.2. **Jack polynomials, Schur polynomials and zonal polynomials.** For some special choices of the deformation parameter α, Jack polynomials coincide (up to some simple normalization constants) with some very established families of symmetric polynomials. In particular, the case $\alpha = 1$ corresponds to *Schur polynomials*, $\alpha = 2$ corresponds to *zonal polynomials* and $\alpha = \frac{1}{2}$ corresponds to *symplectic zonal polynomials*. For these special values of the deformation parameter Jack polynomials are particularly nice because they have some additional structures and features (usually related to algebra and the representation theory) and for this reason they are much better understood.

1.3. **Jack polynomials and maps.** Roughly speaking, a *map* is a graph drawn on a surface, see Figure 1.1. In this article we will investigate the relationship between combinatorics of Jack polynomials and enumeration of maps, related to *generalized characters* and *polynomial functions* on the set of Young diagrams. In the special cases of Schur polynomials ($\alpha = 1$) and zonal polynomials ($\alpha = 2$ and $\alpha = \frac{1}{2}$) this relationship is already well-understood. The generic case is much more mysterious and we will be able only to present some conjectures which generalize these three special cases and — to some extent — to verify positively some consequences of these conjectures.

1.4. **Normalized characters.** The *irreducible character* $\chi^\lambda(\pi) = \operatorname{Tr} \rho^\lambda(\pi)$ of the symmetric group is usually considered as a function of the permutation π, with the Young diagram λ fixed. It was a brilliant observation of

Kerov and Olshanski [KO94] that for several problems of the asymptotic representation theory it is convenient to do the opposite: keep the permutation π fixed and let the Young diagram λ vary. It should be stressed the Young diagram λ is arbitrary, in particular there are no restrictions on the number of boxes of λ. In this way it is possible to study the structure of the series of the symmetric groups $\mathfrak{S}_1 \subset \mathfrak{S}_2 \subset \cdots$ and their representations in a uniform way. This concept is sometimes referred to as *dual combinatorics* of the characters of the symmetric groups.

In order for this idea to be successful one has to replace the usual characters $\chi^\lambda(\pi)$ by, so called, *normalized characters* $\operatorname{Ch}_\pi(\lambda)$, namely for partitions π, λ such that $|\pi| = k$ and $|\lambda| = n$ we define

$$\operatorname{Ch}_\pi(\lambda) = \begin{cases} (n)_k \, \frac{\chi^\lambda(\pi \cup 1^{n-k})}{\chi^\lambda(1^n)} & \text{if } k \leq n, \\ 0 & \text{otherwise,} \end{cases} \tag{1.1}$$

where

$$(n)_k := \underbrace{n(n-1)\cdots(n-i+1)}_{k \text{ factors}}$$

denotes the *falling factorial*.

This choice of normalization is justified by the fact that so defined characters Ch_π belong to the algebra of *polynomial functions on the set of Young diagrams* [KO94], which in the last two decades turned out to be essential for several asymptotic and enumerative problems of the representation theory of the symmetric groups.

1.5. **Jack characters.** Lassalle [Las08, Las09] initiated investigation of a kind of *dual combinatorics* of Jack polynomials. Roughly speaking, it is the investigation of the coefficient standing at $p_{\pi,1,1,\dots}$ in the expansion of Jack symmetric polynomial $J_\lambda^{(\alpha)}$ in the basis of *power-sum symmetric functions*.

The motivation for studying such quantities comes from the observation that in the important special case of Schur polynomials ($\alpha = 1$) one recovers in this way the normalized characters $\operatorname{Ch}_\pi^{(1)} = \operatorname{Ch}_\pi$ given by (1.1) which already proved to have a rich and fascinating structure. The cases of zonal polynomials, symplectic polynomials and general Jack polynomials give rise to some new quantities for which in [FŚ11b] we coined the names *zonal characters* $\operatorname{Ch}_\pi^{(2)}$, *symplectic zonal characters* $\operatorname{Ch}_\pi^{(1/2)}$ and general *Jack characters* $\operatorname{Ch}_\pi^{(\alpha)}$.

Another motivation for studying Jack characters $\operatorname{Ch}_\pi^{(\alpha)}$ comes from the observation that they form a very interesting linear basis of the algebra of α-*polynomial functions on the set of Young diagrams* (which is closely related to the celebrated algebra of polynomial functions).



Figure 1.2. (a) Example of a bipartite graph (drawn on the torus) and (b) an example of its embedding $F(\Sigma) = \alpha$, $F(\Pi) = \beta$, $F(V) = a$, $F(W) = c$. $F(1) = F(4) = (a\beta)$, $F(2) = F(5) = (a\alpha)$, $F(3) = (c\alpha)$. The columns of the Young diagram were indexed by small Latin letters, the rows by small Greek letters.

The main goal of this paper is to *understand the structure of Jack characters* $\mathrm{Ch}_\pi^{(\alpha)}$. In the following we will give more details on this problem.

1.6. **Stanley character formula and embeddings of bipartite graphs.** An *embedding F of a bipartite graph G to a Young diagram λ* is a function which maps the set $V_\circ(G)$ of white vertices of G to the set of columns of λ, which maps the set $V_\bullet(G)$ of black vertices of G to the set of rows of λ, and maps the edges of G to boxes of λ, see Figure 1.2. We also require that an embedding preserves the relation of *incidence*, i.e., a vertex V and an incident edge E should be mapped to a row or column $F(V)$ which contains the box $F(E)$. We denote by $N_G(\lambda)$ the number of such embeddings of G to λ. Quantities $N_G(\lambda)$ were introduced in our paper [FŚ11a] and they proved to be very useful for studying various asymptotic and enumerative problems of the representation theory of symmetric groups [FŚ11a, DFŚ10].

Our main problem of *understanding the structure of Jack characters* $\mathrm{Ch}_\pi^{(\alpha)}$ can be formulated concretely as finding some constants $m_G^{(\alpha)}$ with a property that for any Young diagram λ

$$\mathrm{Ch}_\pi^{(\alpha)}(\lambda) = (-1)^{\ell(\pi)} \sum_G \left(-\frac{1}{\sqrt{\alpha}}\right)^{|V_\bullet(G)|} \left(\sqrt{\alpha}\right)^{|V_\circ(G)|} m_G^{(\alpha)} \, N_G(\lambda), \tag{1.2}$$

where the sum is over some finite family of bipartite graphs. We will refer to this kind of result as *Stanley character formula*. In the following we will discuss the special cases $\alpha \in \left\{\frac{1}{2}, 1, 2\right\}$ for which Stanley character formula is already well-understood and we will discuss some hints about its structure for generic values of the deformation parameter α.

1.7. **Stanley formula for $\alpha = 1$ and oriented maps.** Roughly speaking, *oriented map M* is defined as a bipartite graph drawn on an oriented surface, see Figure 1.1. Since oriented maps are not in the focus of this article, we do not present all necessary definitions and for details we refer to [FŚ11a].

It has been observed in [FŚ11a] that a formula conjectured by Stanley [Sta06, Fér10] for the normalized characters of the symmetric groups can be expressed as the sum

$$\operatorname{Ch}^{(1)}_\pi(\lambda) = \operatorname{Ch}_\pi(\lambda) = (-1)^{\ell(\pi)} \sum_M (-1)^{|V_\bullet(M)|} N_M(\lambda) \tag{1.3}$$

over all *oriented maps M with the face-structure given by π*. Clearly, this formula fits nicely into the template (1.2).

1.8. **Stanley formula for $\alpha = 2, \alpha = 1/2$ and non-oriented maps.** Roughly speaking, a *non-oriented map* is a bipartite graph (possibly with multiple edges) drawn on a surface. For a precise definition we refer to Section 3.1.

In [FŚ11b] it has been observed that

$$\begin{aligned}\operatorname{Ch}^{(2)}_\pi(\lambda) = (-1)^{\ell(\pi)} \sum_M \left(-\frac{1}{\sqrt{2}}\right)^{|V_\bullet(M)|} \left(\sqrt{2}\right)^{|V_\circ(M)|} \times \\ \times \left(-\frac{1}{\sqrt{2}}\right)^{|\pi|+\ell(\pi)-|V(M)|} N_M(\lambda),\end{aligned} \tag{1.4}$$

$$\begin{aligned}\operatorname{Ch}^{(1/2)}_\pi(\lambda) = (-1)^{\ell(\pi)} \sum_M \left(-\sqrt{2}\right)^{|V_\bullet(M)|} \left(\frac{1}{\sqrt{2}}\right)^{|V_\circ(M)|} \times \\ \times \left(\frac{1}{\sqrt{2}}\right)^{|\pi|+\ell(\pi)-|V(M)|} N_M(\lambda),\end{aligned} \tag{1.5}$$

where the sums run over all *non-oriented maps M with the face-type specified by π*. Clearly, also this formula fits nicely into the template (1.2).

1.9. **Orientability generating series and the main conjecture.** As we have seen above, the case $\alpha = 1$ (Eq. (1.3)) corresponds to summation over *oriented maps*, while the cases $\alpha = 2$ and $\alpha = \frac{1}{2}$ correspond to summation over *all* (possibly *non-orientable*) *maps* (Eqs. (1.4), (1.5)) with some simple coefficients which depend only on general features of the map, such as the number of the vertices. Thus one can expect that the summation in Stanley formula (1.2) for the general value of α should involve not summation over some abstract graphs G, but over graphs which carry an additional structure of a *map* M, and the coefficient $m_M^{(\alpha)}$ should be interpreted as a kind of *measure of non-orientability of a given map* M.

This notion of *measure of non-orientability* is not very well defined. For example, one could require that for $\alpha = 1$ the corresponding coefficient $m_M^{(1)}$ is equal to 1 if M is orientable and zero otherwise; and that for $\alpha \in \left\{\frac{1}{2}, 2\right\}$ the coefficient $m_M^{(\alpha)}$ takes some fixed value on all (orientable and non-orientable) maps. This kind of problem already appeared in investigation of the *structure constants* of Jack polynomials [GJ96]; unfortunately it still remains open (for an extensive bibliography on this topic see [LC09]).

In Section 3.5 we will define some quantity mon_M which indeed — although in some perverse sense — measures non-orientability of a given map M. Roughly speaking, it is defined as follows: we remove edges of the map M one after another (in a random order). For each edge which is to be removed we check the *type* of this edge (for example, an edge may be *twisted* if, in some sense, it is a part of *Möbius band*). We multiply the factors corresponding to the types of all edges. The quantity mon_M is defined as the mean value of this product.

This is a rather strange definition. For example, one could complain that this is a weak measure of non-orientability of a map; in particular for $\alpha = 1$ the corresponding weight mon_M does not vanish on non-orientable maps. Nevertheless, we have some indirect evidence that, in fact, mon_M gives the correct answer for the main problem of finding coefficients in Stanley formula for Jack characters. We state it precisely as the following conjecture.

Conjecture 1.1 (The main conjecture)**.** *For arbitrary Young diagram* λ*, partition* π *and* $\alpha > 0$ *the value of Jack character is given by*
(1.6)

$$\operatorname{Ch}_\pi^{(\alpha)}(\lambda) = (-1)^{\ell(\pi)} \sum_M \left(-\frac{1}{\sqrt{\alpha}}\right)^{|V_\bullet(G)|} \left(\sqrt{\alpha}\right)^{|V_\circ(G)|} \operatorname{mon}_M N_M(\lambda),$$

where the summation runs over all non-oriented maps with face-type π*.*

We are unable to prove this conjecture in full generality but in a forthcoming paper [CJDŚ13] we will show several results strongly supporting this conjecture, in particular the following one.

Theorem 1.2 ([CJDŚ13])**.** *Main Conjecture 1.1 is true for partitions* $\pi = (n)$ *consisting of a single part for* $1 \leq n \leq 6$*.*

In the remaining part of this paper we will present some other partial results supporting Main Conjecture 1.1.

We define the *orientability generating series* $\widehat{\mathrm{Ch}}^{(\alpha)}_{\pi}(\lambda)$ as the right hand-side of (1.6):

$$\widehat{\mathrm{Ch}}^{(\alpha)}_{\pi}(\lambda) := (-1)^{\ell(\pi)} \sum_{M} \left(-\frac{1}{\sqrt{\alpha}}\right)^{|V_{\bullet}(M)|} \left(\sqrt{\alpha}\right)^{|V_{\circ}(M)|} \mathrm{mon}_M \ N_M(\lambda), \tag{1.7}$$

where the sum runs over all non-oriented maps M with the face-type π. This expression clearly fits into the template (1.2). Thus, Main Conjecture 1.1 may be equivalently reformulated as follows: for any partition π the corresponding Jack character and the orientability generating series are equal:

$$\mathrm{Ch}^{(\alpha)}_{\pi}(\lambda) = \widehat{\mathrm{Ch}}^{(\alpha)}_{\pi}(\lambda).$$

1.10. **Rectangular Young diagrams.** Investigation of the normalized characters $\mathrm{Ch}_{\pi}(\lambda)$ in the case when $\lambda = p \times q$ is a *rectangular* Young diagram was initiated by Stanley [Sta04] who noticed that they have a particularly simple structure; in particular he showed that formula (1.3) holds true in this special case.

This line of research was continued by Lassalle [Las08] who (apart from other results) studied Jack characters $\mathrm{Ch}^{(\alpha)}_{\pi}(p \times q)$ on rectangular Young diagrams. In particular, Lassalle found a recurrence relation [Las08, formula (6.2)] fulfilled by such characters; this recurrence relates values of the characters on a fixed rectangular Young diagram $p \times q$, corresponding to various partitions π. This recurrence relation is essential for the current paper; our Main Conjecture 1.1 was formulated by a careful attempt of reverse-engineer the hidden hypothetical combinatorial structure behind Lassalle's recurrence.

In particular, our measure of non-orientability of maps mon_M was from the very beginning chosen in such a way that Main Conjecture 1.1 is true for a rectangular Young diagram $\lambda = p \times q$.

We will discuss these issues in Section 5.

1.11. **Consequences of the Main Conjecture.** The indirect evidence which we present in the current paper and rather extensive numerical tests from the forthcoming paper [CJDŚ13] make us confident that Main Conjecture 1.1 is indeed true. For this reason, in Section 6 we dare to explore some of its implications. In particular, Main Conjecture would explain (weaker versions) of some conjectures of Lassalle [Las08].

1.12. **Outlook.**

1.12.1. *Kerov polynomials.* Main Conjecture 1.1 is closely related to *Kerov polynomials*. Since this topic is quite vast, we decided to investigate this relationship in a separate forthcoming paper [CJDŚ13]. In particular, we will discuss there the applications of Main Conjecture 1.1 to conjectures of Lassalle [Las09] concerning Kerov polynomials.

1.12.2. *Polynomial functions on the set of Young diagrams.* The *algebra of polynomial functions on the set of Young diagrams* [KO94] was mentioned only briefly in the current paper. We will discuss its relationship to orientability generating series in a forthcoming paper [Śea13].

1.12.3. *Other problems of combinatorics of Jack polynomials and non-orientability of maps.* In the current paper we investigate the combinatorics of *Jack characters* related to maps. The study of analogous connections between *Jack polynomials* and maps is much older. In particular, Goulden and Jackson [GJ96] formulated a conjecture (called b*-Conjecture*) which claims, roughly speaking, that the connection coefficients of Jack polynomials can be explained combinatorially as summation over certain maps with coefficients which should describe *non-orientability of a given map*. An extensive bibliography to this topic can be found in [LC09].

We conjecture that the measure of non-orientability mon_M which is defined in the current paper provides a *partial* solution to the b-Conjecture of Goulden and Jackson. (We can only claim a *partial* solution because our measure of non-orientability mon_M is a polynomial in variable γ with rational coefficients while Goulden and Jackson were looking for some measure of non-orientability in the form of a single monomial of the form γ^i).

We also conjecture that the problem of finding *structure constants* for Jack characters has a solution given by summation over certain maps M, with the coefficients given by the measure of non-orientability mon_M.

We postpone the precise formulation of our claims for b-Conjecture and the structure constants of Jack characters to forthcoming papers.

2. PRELIMINARIES

2.1. **Partitions and Young diagrams.** A *partition* $\pi = (\pi_1, \dots, \pi_l)$ is defined as a weakly decreasing finite sequence of positive integers. If $\pi_1 + \cdots + \pi_l = n$ we also say that π *is a partition of* n and denote it by $\pi \vdash n$. We will use the following notations: $|\pi| := \pi_1 + \cdots + \pi_l$; furthermore $\ell(\pi) := l$ denotes the *number of parts of* π and

$$m_i(\pi) := \big|\{k : \pi_k = i\}\big|$$

denotes the *multiplicity* of $i \geq 1$ in the partition π.



Figure 2.1. Multirectangular Young diagram $P \times Q$.

When dealing with partitions we will use the shorthand notation

$$1^l := (\underbrace{1, \dots, 1}_{l \text{ times}}).$$

Any partition can be alternatively viewed as a *Young diagram*. For drawing Young diagram we use the French convention.

If $P = (p_1, \dots, p_\ell)$, $Q = (q_1, \dots, q_\ell)$ are sequences of non-negative integers such that $q_1 \geq q_2 \geq \cdots \geq q_\ell$, we consider the *multirectangular* Young diagram

$$P \times Q = (\underbrace{q_1, \dots, q_1}_{p_1 \text{ times}}, \dots, \underbrace{q_\ell, \dots, q_\ell}_{p_\ell \text{ times}}).$$

This concept is illustrated in Figure 2.1. Quantities P and Q are referred to as *Stanley coordinates* of Young diagram $P \times Q$ [Sta06].

The conjugacy classes of the symmetric group $\mathfrak{S}_n$ are in the one-to-one correspondence with partitions of n. Thus any partition $\pi \vdash n$ can be also viewed as some (arbitrarily chosen) permutation $\pi \in \mathfrak{S}_n$ with the corresponding cycle decomposition. This identification between partitions and permutations allows us to define characters such as $\operatorname{Tr} \rho(\pi)$, $\operatorname{Ch}_\pi^{(\alpha)}$ for π being either a permutation or a partition.

2.2. **Jack characters.** *Jack characters* were introduced by Lassalle [Las08, Las09]; since we use a different normalization the following presentation is based on our previous work [DF12].

As there are several of them, we have to fix a normalization for Jack polynomials. In our context, the best is to use the functions denoted by J in the book of Macdonald [Mac95, VI, (10.22)]. We expand Jack polynomial in the basis of power-sum symmetric functions:

$$J_\lambda^{(\alpha)} = \sum_{\substack{\rho: \\ |\rho|=|\lambda|}} \theta_\rho^{(\alpha)}(\lambda)\, p_\rho; \tag{2.1}$$

then we define

$$\mathrm{Ch}_\pi^{(\alpha)}(\lambda) = \alpha^{-\frac{|\pi|-\ell(\pi)}{2}} \binom{|\lambda|-|\pi|+m_1(\pi)}{m_1(\pi)} z_\pi\, \theta^{(\alpha)}_{\pi\cup 1^{|\lambda|-|\pi|}}(\lambda), \tag{2.2}$$

where

$$z_\pi = \pi_1\pi_2\cdots\ m_1(\pi)!\ m_2(\pi)!\cdots .$$

These quantities $\mathrm{Ch}_\pi^{(\alpha)}$ are called *Jack characters*. It turns out that for $\alpha = 1$ we recover the usual normalized characters (1.1) of the symmetric groups.

2.3. **Relationship to Lassalle's normalization.** For reader's convenience we provide below the relationship between quantities used by Lassalle (in boldface) and the ones used by us:

$$\boldsymbol{\vartheta}^\lambda_{\pi\cup 1^{n-|\pi|}}(\alpha) = \binom{|\lambda|-|\pi|+m_1(\pi)}{m_1(\pi)} z_\pi\, \theta^\lambda_{\pi\cup 1^{|\lambda|-|\pi|}}(\alpha),$$

$$\mathrm{Ch}_\pi^{(\alpha)}(\lambda) = \alpha^{-\frac{|\pi|-\ell(\pi)}{2}} \boldsymbol{\vartheta}^\lambda_{\pi\cup 1^{n-|\pi|}}(\alpha). \tag{2.3}$$

3. THE MEASURE OF NON-ORIENTABILITY OF MAPS

3.1. **Non-oriented maps.** In the remaining part of the paper we will consider only *non-oriented bipartite maps*, so we will simplify our notation and we will refer to them as just *maps*. We assume that there are no isolated vertices; multiple edges between vertices are allowed. If we draw an edge of a non-oriented map with a fat pen, its boundary consists of two *edge-sides*. The maps which we consider in this article have labeled edge-sides, see Figure 3.1; in other words each edge carries two labels, one on each of its sides. We also assume that each face of the map is homeomorphic to a disk and each label is unique.

If we unglue each edge of the map into a pair of edge-sides, the map becomes a collection of bipartite polygons, each polygon corresponding to one face of the map, see Figure 3.2. Thus a non-oriented map can be described alternatively as a collection of bipartite polygons with labeled edges (such as the ones from Figure 3.2) together with the pair-partition telling which pairs of edges should be glued together.

Let $2\pi_1, 2\pi_2, \dots, 2\pi_\ell$ be the numbers of edges of the polygons, or — equivalently — the numbers of edges of the faces of the map. We say that $\pi = (\pi_1, \dots, \pi_\ell)$ is the *face-type* of the collection of polygons or the *face-type* of the map.



Figure 3.1. Example of a *non-oriented map* drawn on the projective plane. The left side of the square should be glued with a twist to the right side, as well as bottom to top, as indicated by arrows. The two faces of the map are indicated by the colors.



Figure 3.2. The map from Figure 3.1 is obtained from the above two polygons by gluing the following pairs of edges: $\{1,3\}$, $\{2,10\}$, $\{4,9\}$, $\{5,D\}$, $\{6,C\}$, $\{7,B\}$, $\{8,A\}$. The colors of the polygons correspond to the colors of the faces of the map from Figure 3.1.

Summation over maps with a specified face-type (such as in (1.4) and (1.5)) should be understood as follows: we fix some collection P_π of bipartite polygons with labeled edges of the given face-type π and consider all pair-partitions of their edges; we sum over the resulting collection of



Figure 3.3. The non-oriented map from Figure 3.1. On the boundary of each face some arbitrary orientation was chosen, as indicated by arrows. Edge $\{4, 9\}$ is an example of a straight edge, edge $\{1, 3\}$ is an example of a twisted edge, edge $\{6, C\}$ is an example of an interface edge.



(a) (b) (c)

Figure 3.4. Three possible kinds of edges in a map (see Figure 3.3): (a) *straight edge*: both sides of the edge belong to the same face and have opposite orientations, (b) *twisted edge*: both sides of the edge belong to the same face and have the same orientation, (c) *interface edge*: the sides of the edge belong to two different maps; their orientations are not important. In all three cases the colors of the vertices are not important.

non-oriented maps. The *set of maps with a specified face-type* should be understood in an analogous way.

For a map M let $F(M), E(M), V(M)$ denote for the sets of faces, edges and vertices of M, respectively. For a bipartite map M we recall that a set of vertices of M has a natural decomposition into two sets: the set of white vertices denoted by $V_\circ(M)$ and the set of black vertices denoted by $V_\bullet(M)$.

3.2. **Three kinds of edges.** Let a map M with some selected edge E be given (see example in Figure 3.3). There are three possibilities:

- *Both sides of the edge E are lying on the boundary of the same face F. This means that if we travel along the boundary of the face F then we visit the edge E twice. Assume that the directions in which we travel twice along the edge E are* opposite*, see Figure 3.4a.*

  In this case the edge E is called *straight* and we associate to it the weight 1.
- *Both sides of the edge E are lying on the boundary of the same face F. This means that if we travel along the boundary of the face F then we visit the edge E twice. Assume that the directions in which we travel twice along the edge E are* the same*, see Figure 3.4b.*

  In this case the edge E is called *twisted* and we associate to it the weight $\gamma := \frac{1-\alpha}{\sqrt{\alpha}}$.
- *The edge E is lying on the boundary of two different faces, see Figure 3.4c.*

  In this case the edge E is called *interface* and we associate to it the weight $\frac{1}{2}$.

The weight given by the above convention will be denoted $\operatorname{mon}_{M,E}$.

3.3. **Removal of edges.** If E is an edge of a map M, we denote by $M \setminus \{E\}$ *the map M with the edge E removed*. This definition is a bit subtle; for example since we do now allow maps having isolated vertices, if some endpoint of E is a leaf, we remove it as well. Removal of an edge might change the topology of the surface on which the map is drawn; for this reason instead of figures of the type presented in Figure 3.1 it is more convenient to consider for this purpose *ribbon graphs*, see Figure 3.5.

3.4. **Weight associated to a map with a history.** Let M be a map and let some linear order $\prec$ on the edges be given. This linear order will be called *history*.

Let $E_1, \dots, E_n$ be the sequence of edges of M, listed according to the linear order $\prec$. We set $M_i = M \setminus \{E_1, \dots, E_i\}$ and define

$$\operatorname{mon}_{M,\prec} := \prod_{0 \leq i \leq n-1} \operatorname{mon}_{M_i, E_{i+1}} . \tag{3.1}$$

This quantity can be interpreted as follows: from the map M we remove (one by one) all the edges, in the order specified by the history. For each edge which is about to be removed we consider its weight relative to the current map.



Figure 3.5. Alternative graphical representation of the map from Figure 3.1 as a ribbon graph. The edges of the map are represented as thin ribbons attached to the vertices.

Please note that the type of a given edge (i.e., *straight* versus *twisted* versus *interface*) might change in the process of removing edges and the weight $\operatorname{mon}_{M,\prec}$ usually depends on the choice of the history $\prec$.

3.5. **Measure of non-orientability of a map.** Let M be a map with n edges. We define

$$\operatorname{mon}_M = \operatorname{mon}_M(\gamma) := \frac{1}{n!} \sum_{\prec} \operatorname{mon}_{M,\prec} . \tag{3.2}$$

This quantity can be interpreted as the mean value of the weight associated to the map M equipped with a randomly selected history (with all histories having equal probability). This is the central quantity for the current paper, we call it *the measure of non-orientability* of the map M.

Example 3.1. We consider the map M depicted in Figure 3.6. For the history $A \prec B \prec C$ the corresponding weight is equal to $\operatorname{mon}_{M,\prec} = 1 \cdot \frac{1}{2} \cdot 1$ while for the history $B \prec A \prec C$ the corresponding weight is equal to $\operatorname{mon}_{M,\prec} = \gamma \cdot \gamma \cdot 1$. The other histories are analogous to these two cases,



Figure 3.6. The map M considered in Example 3.1; for simplicity the labels of the edge-sides were removed and each edge carries only one label.

finally

$$\operatorname{mon}_M = \frac{2 \times 1 \cdot \frac{1}{2} \cdot 1 + 4 \times \gamma \cdot \gamma \cdot 1}{6}.$$

Lemma 3.2. *Let M be a map. Then mon_M is a polynomial in variable γ of degree (at most)*

$$d(M) := 2(\textit{number of connected components of } M) - \chi(M),$$

where

$$\chi(M) := |V(M)| - |E(M)| + |F(M)|$$

is the Euler characteristic of M.

The polynomial $\operatorname{mon}_M(\gamma)$ is an even (respectively, odd) polynomial if and only if the Euler characteristic $\chi(M)$ is an even number (respectively, an odd number).

Before we start the proof, notice that if M is a connected, orientable map then $\frac{1}{2}d(M)$ has a natural interpretation as the *genus* of the surface on which M is drawn.

Proof. We claim that for an arbitrary map M and its edge E:

- $\operatorname{mon}_{M,E}$ is a polynomial in γ of degree (at most) $d(M) - d(M \setminus E)$,
- $\operatorname{mon}_{M,E}$ is an even (respectively, odd) polynomial if and only if $d(M) - d(M \setminus E)$ is an even number (respectively, an odd number).

This statement follows by a careful investigation of each of the three cases considered in Section 3.2 (notice that the cases when at least one of the endpoints of E has degree 1 must be considered separately).

From the telescopic product (3.1) it follows that the statement of the lemma holds true if the polynomial mon_M is replaced by $\operatorname{mon}_{M,\prec}$, where $\prec$ is an arbitrary history. The latter result finishes the proof by taking the average over $\prec$. □

4. EVIDENCE FOR THE CONJECTURE: SPECIAL VALUES OF α

Theorem 4.1. *Main Conjecture 1.1 is true for* $\alpha \in \left\{\frac{1}{2}, 2\right\}$.

Proof. Condition $\alpha \in \left\{\frac{1}{2}, 2\right\}$ is equivalent to $\gamma^2 = \frac{1}{2}$. One can check that then for an arbitrary map M and its edge E (case by case analysis, one should separately check the cases when at least one endpoint of E has degree 1)

$$\operatorname{mon}_{M,E} = \gamma^{\left[|F(M)|-|V(M)|)\right]-\left[|F(M\setminus E)|-|V(M\setminus E)|\right]+1}.$$

By a telescopic sum it follows that for an arbitrary history $\prec$:

$$\operatorname{mon}_{M,\prec} = \gamma^{|F(M)|-|V(M)|+|E(M)|} = \gamma^{\ell(\pi)-|V(M)|+|\pi|},$$

where π is the face-type of M; in particular this expression does not depend on the choice of the history.

One can thus easily check that $\widehat{\operatorname{Ch}}^{(\alpha)}_\pi$ coincides with (1.4), respectively (1.5). □

One could expect that the special case of $\alpha = 1$ should be also rather easy, as in this case the Stanley formula (1.3) is already available and $\gamma = 0$ implies that only histories which have no twisted edges might contribute. Unfortunately, we were not able to show that expressions (1.3) and (1.7) coincide in this case.

5. EVIDENCE FOR THE CONJECTURE: RECTANGULAR YOUNG DIAGRAMS AND LASSALLE'S RECURRENCE

The main result of the current section is the following.

Theorem 5.1. *For a rectangular Young diagram*

$$\lambda = p \times q = (\underbrace{q, \dots, q}_{p \text{ times}}),$$

where $p, q \in \mathbb{N}$ *and for arbitrary partition* π *and parameter* $\alpha > 0$, *Main Conjecture 1.1 holds true, i.e.,*

$$\operatorname{Ch}^{(\alpha)}_\pi(\lambda) = \widehat{\operatorname{Ch}}^{(\alpha)}_\pi(\lambda). \tag{5.1}$$

The main idea of the proof is to use recurrence (5.2) found by Lassalle [Las08, formula (6.2)] and to find a combinatorial interpretation of this recurrence.

5.1. **Lassalle's recurrence.** Following Lassalle [Las08] we denote by $\pi \cup (s)$ the partition π with extra part s added and by $\pi \setminus (s)$ the partition π with one part s removed. We also denote

$$\begin{aligned} \pi_{\uparrow(s)} &= \pi \setminus (s) \cup (s+1), & \pi_{\downarrow(s)} &= \pi \setminus (s) \cup (s-1), \\ \pi_{\uparrow(rs)} &= \pi \setminus (r+s+1) \cup (r,s), & \pi_{\downarrow(rs)} &= \pi \setminus (r,s) \cup (r+s-1), \\ \pi \cup 1^l &= \pi \cup \underbrace{(1) \cup \cdots \cup (1)}_{l \text{ times}}. \end{aligned}$$

Consider a rectangular Young diagram $\lambda = p \times q$ and a partition π such that $m_1(\pi) = 0$ (i.e., π does not contain any part equal to 1). Then Lassalle's recurrence relation [Las08, formula (6.2)], after adapting to our normalizations takes the form:

$$\begin{aligned} (5.2) \quad & \underbrace{\left(\frac{p}{\sqrt{\alpha}} - \sqrt{\alpha} q\right) \sum_r r\, m_r(\pi)\ \mathrm{Ch}^{(\alpha)}_{\pi_{\downarrow(r)}}(\lambda)}_{\text{(removing a leaf)}} \\ & + \underbrace{\sum_r r\, m_r(\pi) \sum_{i=1}^{r-2} \mathrm{Ch}^{(\alpha)}_{\pi_{\uparrow(i,r-i-1)}}(\lambda)}_{\text{(removing a straight edge)}} \\ & - \underbrace{\gamma \sum_r r(r-1)\, m_r(\pi)\ \mathrm{Ch}^{(\alpha)}_{\pi_{\downarrow(r)}}(\lambda)}_{\text{(removing a twisted edge)}} \\ & + \underbrace{\sum_{r,s} rs\, m_r(\pi)\big(m_s(\pi) - \delta_{r,s}\big)\ \mathrm{Ch}^{(\alpha)}_{\pi_{\downarrow(rs)}}(\lambda)}_{\text{(removing an interface edge)}} \\ & = -|\pi|\ \mathrm{Ch}^{(\alpha)}_{\pi}(\lambda). \end{aligned}$$

The comments concerning individual summands on the left-hand side of this recurrence relation are connected to our combinatorial interpretation of this formula and will be explained later on.

The main difficulty with this formula comes from the fact that it was proved only under assumption that $m_1(\pi) = 0$. In the following we will show how to overcome this issue.

5.2. **Partitions with parts equal to** 1**.** In the case $\alpha = 1$, Jack characters $\mathrm{Ch}^{(1)}_\pi = \mathrm{Ch}_\pi$ are simply characters of the symmetric groups and the partition π can be interpreted as the conjugacy class on which we evaluate the character, see (1.1). Thus the parts of π which are equal to 1 correspond to fix-points. Such fix-points are trivial in the sense that do not change the value of the non-normalized character $\operatorname{Tr} \rho(\pi)$; in the case of the normalized characters Ch_π they change the value of the character in a very simple way, namely the identity

$$\mathrm{Ch}^{(\alpha)}_{\pi\cup 1^l}(\lambda) = (|\lambda| - |\pi|)_l \ \mathrm{Ch}^{(\alpha)}_\pi(\lambda) \tag{5.3}$$

holds true for $\alpha = 1$. Strictly from the definition of Jack characters (2.1), (2.2) it follows that the above identity holds true for general value of $\alpha > 0$ and arbitrary Young diagram λ.

The following result shows that analogous property is fulfilled by $\widehat{\mathrm{Ch}}^{(\alpha)}_\pi$ as well.

Lemma 5.2. *Let π be a partition and λ be an arbitrary Young diagram. Then*

$$\widehat{\mathrm{Ch}}^{(\alpha)}_{\pi\cup 1^l}(\lambda) = (|\lambda| - |\pi|)_l \, \widehat{\mathrm{Ch}}^{(\alpha)}_\pi(\lambda). \tag{5.4}$$

Proof. It is enough to prove that for any partition π the following holds:

$$\widehat{\mathrm{Ch}}^{(\alpha)}_{\pi\cup(1)}(\lambda) = (|\lambda| - |\pi|) \ \widehat{\mathrm{Ch}}^{(\alpha)}_\pi(\lambda). \tag{5.5}$$

For a partition π let F_π be the set of pairs $(M, \prec)$, where M is a map with a face-type π and with history $\prec$. More explicitly: we start by fixing some family P_π of polygons of face-type π; then there is a bijective correspondence between maps of face-type π and pair-partitions of the edges of the polygons from P_π. In other words, each element of F_π can be viewed as a pair-partition on edges of P_π, equipped with some linear order on the pairs.

With these notations, a map M with face-type $\pi \cup (1)$ can be viewed as a pair-partition of the edges of the polygons from the family $P_{\pi\cup(1)}$. We have freedom of choosing the family of polygons, therefore we can set $P_{\pi\cup(1)} := P_\pi \sqcup B$, where B is a marked *bigon* (bipartite polygon with two vertices connected by edges denoted b_1, b_2).

If $(M, \prec) \in F_{\pi\cup(1)}$ is such that $M \ni \{b_1, b_2\}$ is a pair-partition which connects the edge b_1 with b_2, we say that $(M, \prec) \in F^0_{\pi\cup(1)}$; in the other case we say that $(M, \prec) \in F^1_{\pi\cup(1)}$. This gives a disjoint decomposition

$$F_{\pi\cup(1)} = F^0_{\pi\cup(1)} \sqcup F^1_{\pi\cup(1)}. \tag{5.6}$$

Let $\mathcal{H} : F_{\pi\cup(1)} \to F_\pi$ be a function $\mathcal{H} : (M, \prec) \mapsto (M', \prec')$ defined as follows:

- If $(M,\prec) \in F^0_{\pi\cup(1)}$ then $M' := M \setminus \big\{\{b_1,b_2\}\big\}$ is defined as the pair-partition M with the pair formed by the edges of B removed; as the linear order $\prec'$ we take the restriction of $\prec$ to $M' \subset M$.

  In this case M, viewed as a bipartite graph, is a disjoint sum of the bipartite graph M' and the bipartite graph consisting of two vertices connected by the edge $\{b_1,b_2\}$. The process of calculating $\operatorname{mon}_{M,\prec}$ is almost identical to the analogous process of calculating $\operatorname{mon}_{M',\prec'}$ except for the additional edge $\{b_1,b_2\}$ which is clearly a straight edge. Thus

$$\begin{aligned} N_M(\lambda) =& |\lambda|\ N_{M'}(\lambda), \\ |V_\circ(M)| =& |V_\circ(M')|+1, \\ |V_\bullet(M)| =& |V_\bullet(M')|+1, \\ \operatorname{mon}_{M,\prec} =& \operatorname{mon}_{M',\prec'}. \end{aligned}$$

- If $(M,\prec) \in F^1_{\pi\cup(1)}$ then there are two pairs of edges $\{e_1,b_i\},\{e_2,b_j\} \in M$ which connect the edges of the marked bigon B with some other edges; we choose the indices i,j in such a way that $\{i,j\}=\{1,2\}$ and $\{e_1,b_i\} \prec \{e_2,b_j\}$. Then

$$M' := \big(M \cup \big\{\{e_1,e_2\}\big\}\big) \setminus \big\{\{e_1,b_i\},\{e_2,b_j\}\big\}$$

  is defined as the map obtained from M by ungluing the two edges glued to the bigon B, removing the bigon B and gluing the two unglued edge-sides together. As the linear order $\prec'$ we take the unique linear order which coincides with $\prec$ on the intersection of their domains and such that for any pair $P \in M' \cap M$ we have that

$$P \prec' \{e_1,e_2\} \iff P \prec \{e_2,b_2\};$$

  in other words the order $\prec'$ is obtained from $\prec$ by substituting the pair $\{e_2,b_2\}$ by $\{e_1,e_2\}$.

  The map M is obtained from M' by replacing the edge $\{e_1,e_2\}$ by a pair of edges in such a way that a new face is created (this is the face corresponding to the bigon B). The process of calculating $\operatorname{mon}_{M,\prec}$ is almost identical to the analogous process of calculating $\operatorname{mon}_{M',\prec'}$ except for the edge $\{e_1,b_1\}$ which is the one of the two edges adjacent to the bigon B which is removed first. This edge is clearly an interface edge. Thus

$$\begin{aligned} N_M(\lambda) =& N_{M'}(\lambda), \\ |V_\circ(M)| =& |V_\circ(M')|, \\ |V_\bullet(M)| =& |V_\bullet(M'), \\ \operatorname{mon}_{M,\prec} =& \frac{1}{2} \operatorname{mon}_{M',\prec'}. \end{aligned}$$

The left-hand side of (5.5) is equal to

$$\begin{aligned} &\widehat{\operatorname{Ch}}^{(\alpha)}_{\pi\cup(1)}(\lambda) \\ &= \sum_{(M,\prec)\in F_{\pi\cup(1)}} \frac{(-1)^{\ell(\pi)+1}}{(|\pi|+1)!} \left(-\frac{1}{\sqrt{\alpha}}\right)^{|V_\bullet(M)|} \left(\sqrt{\alpha}\right)^{|V_\circ(M)|} \operatorname{mon}_{M,\prec}\ N_M(\lambda) \\ &= \sum_{(M,\prec)\in F^0_{\pi\cup(1)}} \frac{(-1)^{\ell(\pi)}}{(|\pi|+1)!} \left(-\frac{1}{\sqrt{\alpha}}\right)^{|V_\bullet(M')|} \left(\sqrt{\alpha}\right)^{|V_\circ(M')|} \operatorname{mon}_{M',\prec'} |\lambda|\, N_{M'}(\lambda) \\ &- \sum_{(M,\prec)\in F^1_{\pi\cup(1)}} \frac{(-1)^{\ell(\pi)}}{(|\pi|+1)!} \left(-\frac{1}{\sqrt{\alpha}}\right)^{|V_\bullet(M')|} \left(\sqrt{\alpha}\right)^{|V_\circ(M')|} \operatorname{mon}_{M',\prec'} \frac{1}{2}\, N_{M'}(\lambda), \end{aligned} \tag{5.7}$$

where $\mathcal{H} : (M,\prec) \mapsto (M',\prec')$ and in the last equality we used the decomposition (5.6).

In the following we will show that for each $(M',\prec') \in F_\pi$ its preimage fulfills:

$$\begin{aligned} |\mathcal{H}^{-1}(M',\prec') \cap F^0_{\pi\cup(1)}| &= |\pi|+1, \\ |\mathcal{H}^{-1}(M',\prec') \cap F^1_{\pi\cup(1)}| &= 2(|\pi|+1)|\pi|. \end{aligned}$$

Indeed, for all $(M,\prec) \in \mathcal{H}^{-1}(M',\prec') \cap F^0_{\pi\cup(1)}$ the maps $M = M' \cup \big\{\{b_1,b_2\}\big\}$ are all the same; the order $\prec$ is obtained from the order $\prec'$ by adding the additional pair $\{b_1,b_2\}$ between pairs forming M' and this can be done in $|\pi|+1$ ways.

Similarly, $(M,\prec) \in \mathcal{H}^{-1}(M',\prec') \cap F^1_{\pi\cup(1)}$ can be obtained from $(M',\prec')$ by removing some pair $\{e_1,e_2\} \in M'$, and adding pairs $\{e_1,b_i\}$ and $\{e_2,b_j\}$ for some choice of $\{i,j\} = \{1,2\}$. Since we can replace the roles of the edges e_1 and e_2, we have altogether 4 choices for doing this. The pair $\{e_1,e_2\} \in M'$ can be equivalently specified by saying that there are ℓ elements which are smaller than $\{e_1,e_2\}$ (with respect to $\prec'$). The linear order $\prec$ is obtained by substituting the pair $\{e_1,e_2\}$ by $\{e_2,b_j\}$ and by adding the pair $\{e_1,b_i\}$ in such a way that $\{e_1,b_i\} \prec \{e_2,b_j\}$; there are $\ell+1$ choices for this. Thus the total number of choices is equal to

$$\sum_{0\leq \ell \leq |\pi|-1} 4(\ell+1) = 2(|\pi|+1)|\pi|,$$

just as we claimed.

Concluding, it gives us

$$\begin{aligned}
&\widehat{\operatorname{Ch}}^{(\alpha)}_{\pi\cup(1)}(\lambda) \\
&= \frac{1}{|\pi|+1} \sum_{(M',\prec')\in F_\pi} \frac{(-1)^{\ell(\pi)}}{(|\pi|)!} \left(-\frac{1}{\sqrt{\alpha}}\right)^{|V_\bullet(M')|} \left(\sqrt{\alpha}\right)^{|V_\circ(M')|} \\
&\qquad\qquad \times \operatorname{mon}_{M',\prec'}\ N_{M'}(\lambda) \left(|\lambda|(|\pi|+1) - 2(|\pi|+1)|\pi|\frac{1}{2}\right) \\
&= (|\lambda|-|\pi|) \\
&\times \sum_{(M',\prec')\in F_\pi} \frac{(-1)^{\ell(\pi)}}{(|\pi|)!} \left(-\frac{1}{\sqrt{\alpha}}\right)^{|V_\bullet(M')|} \left(\sqrt{\alpha}\right)^{|V_\circ(M')|} \operatorname{mon}_{M',\prec'}\ N_{M'}(\lambda) \\
&\qquad\qquad\qquad\qquad = (|\lambda|-|\pi|)\, \widehat{\operatorname{Ch}}^{(\alpha)}_{\pi}(\lambda)
\end{aligned}$$

which finishes the proof. □

5.3. **Recurrence relation for the orientability generating series.** The following result states that the orientability generating series $\widehat{\operatorname{Ch}}^{(\alpha)}$ fulfill the recurrence relation analogous to (5.2).

Proposition 5.3. *If $\lambda = p \times q$ is a rectangular Young diagram and π is a partition such that $m_1(\pi) = 0$ then*

$$\begin{aligned} &\underbrace{\left(\frac{p}{\sqrt{\alpha}} - \sqrt{\alpha} q\right) \sum_r r\, m_r(\pi)\, \widehat{\mathrm{Ch}}^{(\alpha)}_{\pi\downarrow(r)}(\lambda)}_{\text{(removing a leaf)}} \\ &\qquad + \underbrace{\sum_r r\, m_r(\pi) \sum_{i=1}^{r-2} \widehat{\mathrm{Ch}}^{(\alpha)}_{\pi\uparrow(i,r-i-1)}(\lambda)}_{\text{(removing a straight edge)}} \\ &\qquad - \underbrace{\gamma \sum_r r(r-1)\, m_r(\pi)\, \widehat{\mathrm{Ch}}^{(\alpha)}_{\pi\downarrow(r)}(\lambda)}_{\text{(removing a twisted edge)}} \\ &\qquad + \underbrace{\sum_{r,s} rs\, m_r(\pi)\big(m_s(\pi) - \delta_{r,s}\big)\, \widehat{\mathrm{Ch}}^{(\alpha)}_{\pi\downarrow(rs)}(\lambda)}_{\text{(removing an interface edge)}} \\ &\qquad\qquad = -|\pi|\, \widehat{\mathrm{Ch}}^{(\alpha)}_{\pi}(\lambda). \end{aligned} \tag{5.8}$$

Proof. The right-hand side of (5.8) is equal to

$$\begin{aligned} &-|\pi|\, \widehat{\mathrm{Ch}}^{(\alpha)}_{\pi}(\lambda) \\ &\qquad = \frac{(-1)^{\ell(\pi)-1}}{(|\pi|-1)!} \sum_{(M,\prec)} \left(-\frac{p}{\sqrt{\alpha}}\right)^{|V_\bullet(M)|} \big(q\sqrt{\alpha}\big)^{|V_\circ(M)|} \mathrm{mon}_{M,\prec}, \end{aligned} \tag{5.9}$$

where the summation is over M with face-type π. We used here that for a rectangular Young diagram $\lambda = p \times q$, the number of embeddings

$$N_M(\lambda) = p^{|V_\bullet(M)|}\, q^{|V_\circ(M)|}$$

takes a particularly simple form. Let

$$\mathrm{contribution}_{M,\prec}(\lambda) := \left(-\frac{p}{\sqrt{\alpha}}\right)^{|V_\bullet(M)|} \big(q\sqrt{\alpha}\big)^{|V_\circ(M)|} \mathrm{mon}_{M,\prec}$$

denote the contribution of the pair $(M, \prec)$ to the right-hand side of (5.9).

Let E be the first edge of M, according to the linear order $\prec$ and let $\prec'$ denote the restriction of the linear order $\prec$ to the edges of $M \setminus E$. Clearly, (3.1) is equivalent to a recursive relationship

$$\mathrm{mon}_{M,\prec} = \mathrm{mon}_{M,E} \cdot \mathrm{mon}_{M\setminus E,\prec'}\,.$$

The summation over $\prec$ can be rewritten as a summation over all choices for E and over all choices of $\prec'$ (linear orders on the edges of $M \setminus E$).

In the following we will show how this allows to rewrite the right-hand side of (5.9). According to the classification from Section 3.2 there are the following possibilities:

- The edge E is straight and thus $\operatorname{mon}_{M,E} = 1$. There are the following three possibilities:
  - Both endpoints of E have degree 1. This is not possible since it would imply that one of the faces of M is a bigon thus $m_1(\pi) \geq 1$.
  - Only the black (respectively, white) endpoint of E has degree 1 (i.e., it is a leaf). We consider the map $M \setminus E$; since we do not allow maps with vertices of degree 0, together with the edge E we also remove its black (respectively, white) endpoint. It follows that

    $$\operatorname{contribution}_{M,\prec}(\lambda) = \frac{-p}{\sqrt{\alpha}} \operatorname{contribution}_{M\setminus E,\prec'}(\lambda);$$

    respectively,

    $$\operatorname{contribution}_{M,\prec}(\lambda) = q\,\sqrt{\alpha}\, \operatorname{contribution}_{M\setminus E,\prec'}(\lambda).$$

    The face-type of $M \setminus E$ is equal to $\pi \downarrow (r)$ with $r \geq 2$, where $2r$ is the number of the edges of the face of M which is adjacent to E.

    The summation over (M, E) such as considered in the current case can be alternatively performed as follows. As usual, let P_π be a collection of bipartite polygons with face-type π. We select some polygon P from the family P_π in such a way that P has $2r$ edges. We choose a pair of consecutive edges of P sharing a black (respectively, white) vertex, we glue together these edges (they form the edge E) and we remove these edges from P. The resulting family of polygons has face-type $\pi \downarrow (r)$; we denote it by $P_{\pi\downarrow(r)}$.

    Polygon P and its two consecutive edges can be selected in $m_r(\pi)\, r$ ways. Then we consider all maps M' resulting from gluing the edges of $P_{\pi\downarrow(r)}$. The multiset consisting of M' given by this construction coincides with the multiset consisting of $M \setminus E$ given by the previous construction. Thus the total contribution to the right-hand side of (5.9) coming from $(M, \prec)$

such as considered in the current case is equal to

$$\sum_r r\, m_r(\pi) \sum_{M',\prec'} \frac{(-1)^{\ell(\pi)}}{(|\pi|-1)!} \left(\frac{p}{\sqrt{\alpha}} - \sqrt{\alpha} q \right) \operatorname{contribution}_{M',\prec'}(\lambda) \\ = \left(\frac{p}{\sqrt{\alpha}} - \sqrt{\alpha} q \right) \sum_r r\, m_r(\pi)\, \widehat{\operatorname{Ch}}_{\pi\downarrow(r)}(\lambda),$$

where the summation on the left-hand side is over maps M' with face-type $\pi \downarrow (r)$. Clearly, it is equal to the first summand on the left-hand side of (5.8).

- No endpoint of E has degree 1; then

  $$\operatorname{contribution}_{M,\prec}(\lambda) = \operatorname{contribution}_{M\setminus E,\prec'}(\lambda).$$

  After removing the edge E, the face of M which was adjacent to E is split into two faces of $M \setminus E$: F_1 and F_2. We use the convention that F_1 (respectively, F_2) is the face which contains the black (respectively, the white) endpoint of the edge E. We denote the number of edges of F_1 by $2i$ and the number of edges of F_2 by $2j$. In other words, the face-type of $M \setminus E$ is equal to $\pi \uparrow (i,j)$ for some $i, j \geq 1$.

  The summation over (M, E) such as considered in the current case can be alternatively performed as follows. We select some polygon P from the family P_π such that P has $2r$ edges. We choose a pair of non-consecutive edges of P such that they are separated by an *even* number of edges, we glue them together (they form the edge E) and we remove them. In this way, polygon P splits into two polygons; we denote by $2i$ (respectively, $2j$) the number of the edges of the polygon which contains the black (respectively, the white) endpoint of E. We have $r = i + j + 1$. The resulting family of polygons has face-type $\pi \uparrow (i,j)$ thus can be identified with $P_{\pi\uparrow(i,j)}$.

  For a fixed choice of $i, j \geq 1$, the polygon P and the pair of edges can be selected in $r\, m_r(\pi)$ ways. Then we consider all maps M' resulting from gluing edges of $P_{\pi\uparrow(i,j)}$. The multiset containing M' given by this construction coincides with the multiset containing $M \setminus E$ given by the previous construction. Thus the total contribution to the right-hand side of (5.9) coming from $(M, \prec)$ such as considered in the current case is equal to

$$\sum_r r\, m_r(\pi) \sum_{i=1}^{r-2} \sum_{M',\prec'} \frac{(-1)^{\ell(\pi\uparrow(i,r-i-1))}}{(k-1)!} \operatorname{contribution}_{M',\prec'}(\lambda) \\ = \sum_r r\, m_r(\pi) \sum_{i=1}^{r-2} \operatorname{Ch}^{(\alpha)}_{\pi\uparrow(i,r-i-1)}(\lambda)$$

where the summation on the left-hand side is over maps M' with face-type $\pi \uparrow (i, r-i-1)$. Clearly, it is equal to the second summand on the left-hand side of (5.8).

- The edge E is twisted and thus $\operatorname{mon}_{M,E} = \gamma$. Then, no endpoint of E has degree 1, hence

$$\operatorname{contribution}_{M,\prec}(\lambda) = \gamma \operatorname{contribution}_{M\setminus E,\prec'}(\lambda).$$

The face-type of $M \setminus E$ is equal to $\pi \downarrow (r)$ with $r \geq 2$, where $2r$ is the number of the edges of the face of M which is adjacent to E.

The summation over (M, E) such as considered in the current case can be alternatively performed as follows. We select some polygon P from the family P_π such that P has $2r$ edges. We choose a pair of edges such that there is an *odd* number of edges between them, we glue them (they form the edge E) and we remove them. The resulting family of polygons has face-type $\pi \downarrow (r)$ thus can be identified with $P_{\pi\downarrow(r)}$.

Polygon P and its two consecutive edges can be selected in $m_r(\pi)\, r(r-1)$ ways. Then we consider all maps M' resulting from gluing edges of $P_{\pi\downarrow(r)}$. The multiset containing M' given by this construction coincides with the multiset consisting of $M \setminus E$ given by the previous construction. Thus the total contribution to the right-hand side of (5.9) coming from $(M, \prec)$ such as considered in the current case is equal to

$$\sum_r r(r-1)\, m_r(\pi) \sum_{M',\prec'} \frac{(-1)^{\ell(\pi)}}{(|\pi|-1)!} \gamma \operatorname{contribution}_{M',\prec'}(\lambda) \\ = \gamma \sum_r r(r-1)\, m_r(\pi)\, \widehat{\operatorname{Ch}}_{\pi\downarrow(r)}(\lambda),$$

where the summation on the left-hand side is over maps M' with face-type $\pi \downarrow (r)$. Clearly, it is equal to the third summand on the left-hand side of (5.8).

- The edge E is interface and thus $\operatorname{mon}_{M,E} = \frac{1}{2}$. Then, no endpoint of E has degree 1, hence

  $$\begin{aligned} N_M(\lambda) =& N_{M\setminus E}(\lambda), \\ \operatorname{contribution}_{M,\prec}(\lambda) =& \frac{1}{2} \operatorname{contribution}_{M\setminus E,\prec'}(\lambda). \end{aligned}$$

  The face-type of $M \setminus E$ is equal to $\pi \downarrow (rs)$ where $2r$ and $2s$ are the numbers of the edges of the faces of M which are adjacent to E.

  The summation over (M, E) such as considered in the current case can be alternatively performed as follows. We select two polygons P_1, P_2 from the family P_π such that P_1 has $2r$ edges and P_2 has $2s$ edges. We choose one edge from P_1 and one edge from P_2, we glue them (they form the edge E) and we remove them. The resulting family of polygons has face-type $\pi \downarrow (rs)$ thus can be identified with $P_{\pi\downarrow(rs)}$.

  Polygons P_1, P_2 can be selected in $m_r(\pi)(m_s(\pi) - \delta_{rs})$ ways and the pair of edges can be selected in $4rs$ ways. Since the roles of P_1 and P_2 can be interchanged, there is an additional factor $\frac{1}{2}$. Then we consider all maps M' resulting from gluing the edges of $P_{\pi\downarrow(rs)}$. The multiset containing M' given by this construction coincides with the multiset consisting of $M \setminus E$ given by the previous construction. Thus the total contribution to the right-hand side of (5.9) coming from $(M, \prec)$ such as considered in the current case is equal to

  $$\begin{aligned} \sum_{r,s} 2rs\, m_r(\pi)\big(m_s(\pi) - \delta_{r,s}\big) \sum_{M',\prec'} \frac{(-1)^{\ell(\pi)}}{(|\pi|-1)!} \frac{1}{2} \operatorname{contribution}_{M',\prec'}(\lambda) \\ = \sum_{r,s} rs\, m_r(\pi)\big(m_s(\pi) - \delta_{r,s}\big)\ \widehat{\operatorname{Ch}}_{\pi\downarrow(r)}(\lambda), \end{aligned}$$

  where the summation on the left-hand side is over maps M' with face-type $\pi \downarrow (r)$. Clearly, it is equal to the third summand on the left-hand side of (5.8).

□

5.4. **Proof of Theorem 5.1.**

Proof of Theorem 5.1. We will use induction over $|\pi|$. For $|\pi| = 0$ there is only the empty partition $\pi = \emptyset$; clearly in this case $\operatorname{Ch}^{(\alpha)}_{\emptyset}(\lambda) = \widehat{\operatorname{Ch}}^{(\alpha)}_{\emptyset}(\lambda) = 1$ holds true. Since this is a bit pathological case (empty polygon, empty function, etc), in order to avoid difficulties with the start of the induction, we also consider separately the case $|\pi| = 1$ for which there is only one partition $\pi = (1)$; we easily get that that $\mathrm{Ch}_1^{(\alpha)}(\lambda) = \widehat{\mathrm{Ch}}_1^{(\alpha)}(\lambda) = |\lambda|$ indeed holds true.

Let us assume that the inductive assertion holds for all π such that $|\pi| < n$ and let π be a partition with $|\pi| = n$. In the case when $m_1(\pi) \geq 1$ we apply (5.3) and (5.4) and the inductive assertion implies that (5.1) holds true for π as well.

In the case when $m_1(\pi) = 0$, we compare the left-hand side of (5.8) with the left-hand side of (5.2). From the inductive assertion it follows that they are equal; so must be their right-hand sides. □

6. CONSEQUENCES OF THE MAIN CONJECTURE: POSITIVITY CONJECTURES OF LASSALLE

The main motivation for our conjectural formula (1.6) for Jack characters was an attempt to prove two conjectures of Lassalle [Las08, Las09]. Both of them state some kind of positivity results, which suggests existence of the underlying combinatorial structure describing Jack characters. Moreover, these conjectures generalize two very interesting ex-conjectures describing structures of the normalized characters of the symmetric group, namely Stanley formula [Sta06] and Kerov conjecture [Ker00, Bia03]; in fact these two results correspond to the substitution $\alpha = 1$ in Lassalle's conjectures.

We postpone the discussion of results and conjectures related to Kerov polynomials and free cumulants to a forthcoming paper [CJDŚ13]. In this section we are going to describe the conjecture of Lassalle [Las08] which concerns Stanley's coordinates and discuss its relation with Main Conjecture 1.1.

6.1. Positivity in Stanley's coordinates.

In the paper [Las08] Lassalle stated the following conjecture.

Conjecture 6.1 ([Las08, Conjecture 1])**.** *Let π be a partition such that $m_1(\pi) = 0$ and let $\beta := \alpha - 1$. Then $(-1)^{|\pi|}\boldsymbol{\vartheta}^{P\times Q}_{\pi\cup 1^{n-|\pi|}}(\alpha)$ is a polynomial in $(P, -Q, \beta)$ with non-negative integer coefficients. These coefficients have a combinatorial interpretation.*

In the following we will prove (in Corollary 6.3) a weaker version of this conjecture. Our main tool will be the following result.

Theorem 6.2. *Let us assume that Main Conjecture 1.1 holds true. Then $(-1)^{|\pi|}\,\mathrm{Ch}_\pi^{(\alpha)}(P \times Q)$ is a polynomial in variables $(P/\sqrt{\alpha}, -\sqrt{\alpha}Q, -\gamma)$ with non-negative rational coefficients.*

Proof. For a Young diagram $P \times Q$ given in the Stanley coordinates, the number of embeddings $N_M(P \times Q)$ takes a particularly simple form (see

[FŚ11b, Lemma 3.9] where the notation is slightly different), for this reason (1.6) would imply that

$$\mathrm{Ch}_{\pi}^{(\alpha)}(P \times Q) = (-1)^{\ell(\pi)} \sum_{M} \mathrm{mon}_M(\gamma) \times \\ \left[\sum_{\varphi: V_\bullet(M) \to \mathbb{N}^\star} \prod_{l \in V_\bullet(M)} \left(\frac{-p_{\varphi(l)}}{\sqrt{\alpha}} \right) \cdot \prod_{l' \in V_\circ(M)} \left(\sqrt{\alpha}\, q_{\psi(l')} \right) \right],$$

where the first sum is over all bipartite maps M of the face-type π and $\psi(l')$ is defined as the maximum of $\varphi(l)$ over all white neighbors l of the black vertex l.

Quantity $\mathrm{mon}_M(\gamma)$ is a polynomial in γ with non-negative rational coefficients. Furthermore, Lemma 3.2 shows that

$$(-1)^{|\pi|} \mathrm{Ch}_{\pi}^{(\alpha)}(P \times Q) = \sum_{M} \mathrm{mon}_M(-\gamma) \times \\ \left[\sum_{\varphi: V_\bullet(M) \to \mathbb{N}^\star} \prod_{l \in V_\bullet(M)} \left(\frac{p_{\varphi(l)}}{\sqrt{\alpha}} \right) \cdot \prod_{l' \in V_\circ(M)} \left(-\sqrt{\alpha} q_{\psi(l')} \right) \right],$$

which finishes the proof. □

Corollary 6.3. *Let us assume that Main Conjecture 1.1 holds true. Let* π *be an arbitrary partition. Then* $(-1)^{|\pi|} \boldsymbol{\vartheta}^{P \times Q}_{\pi \cup 1^{n-|\pi|}}(\alpha)$ *is a polynomial in* $(P, -Q, \beta)$ *with non-negative rational coefficients.*

Proof. Substituting (2.3) we obtain:

$$(-1)^{|\pi|} \boldsymbol{\vartheta}^{P \times Q}_{\pi \cup 1^{n-|\pi|}}(\alpha) = \sum_{M} \sqrt{\alpha}^{2|V_\circ(M)| + |\pi| - \ell(\pi) - |V(M)|} \mathrm{mon}_M(-\gamma) \\ \times \left[\sum_{\varphi: V_\bullet(M) \to \mathbb{N}^\star} \prod_{l \in V_\bullet(M)} p_{\varphi(l)} \cdot \prod_{l' \in V_\circ(M)} \left(-q_{\psi(l')} \right) \right].$$

By Lemma 3.2, $\mathrm{mon}_M(\gamma)$ is a polynomial of degree at most

$$2(\text{number of connected components of } M) - \ell(\pi) + |\pi| - |V(M)|$$

and with the same parity as

$$2(\text{number of connected components of } M) - \ell(\pi) + |\pi| - |V(M)|.$$

The number of connected components of M is at most equal to the number of white vertices, and since $-\gamma = \frac{\beta}{\sqrt{\alpha}}$ and $\alpha = \beta + 1$ we have that

$$\sqrt{\alpha}^{2|V_\circ(M)| + |\pi| - \ell(\pi) - |V(M)|} \mathrm{mon}_M(-\gamma)$$

is a polynomial in β with non-negative rational coefficients. It finishes the proof. □

ACKNOWLEDGMENTS

M.D.'s research has been supported by a grant of *Narodowe Centrum Nauki* (2011/03/N/ST1/00117). In the initial phase of research, P.Ś. was a holder of a fellowship of *Alexander von Humboldt-Stiftung*. P.Ś.'s research has been supported by a grant of *Deutsche Forschungsgemeinschaft* (SN 101/1-1).

REFERENCES

[Bia03] Philippe Biane. Characters of symmetric groups and free cumulants. In *Asymptotic combinatorics with applications to mathematical physics (St. Petersburg, 2001)*, volume 1815 of *Lecture Notes in Math.*, pages 185–200. Springer, Berlin, 2003.

[CJDŚ13] Agnieszka Czyżewska-Jankowska, Maciej Dołęga, and Piotr Śniady. Jack polynomials and the measure of non-orientability of maps. Part 2. Kerov polynomials and genus expansion. In preparation, 2013.

[DF12] Maciej Dołęga and Valentin Féray. On Kerov polynomials for Jack characters. Preprint arXiv:1201.1806v1, 2012.

[DFŚ10] Maciej Dołęga, Valentin Féray, and Piotr Śniady. Explicit combinatorial interpretation of Kerov character polynomials as numbers of permutation factorizations. *Adv. Math.*, 225(1):81–120, 2010.

[Fér10] V. Féray. Stanley's formula for characters of the symmetric group. *Annals of Combinatorics*, 13(4):453–461, 2010.

[FŚ11a] Valentin Féray and Piotr Śniady. Asymptotics of characters of symmetric groups related to Stanley character formula. *Ann. of Math. (2)*, 173(2):887–906, 2011.

[FŚ11b] Valentin Féray and Piotr Śniady. Zonal polynomials via Stanley's coordinates and free cumulants. *J. Algebra*, 334:338–373, 2011.

[GJ96] I. P. Goulden and D. M. Jackson. Connection coefficients, matchings, maps and combinatorial conjectures for Jack symmetric functions. *Trans. Amer. Math. Soc.*, 348(3):873–892, 1996.

[GR05] Adriano Garsia and Jeffrey B. Remmel. Breakthroughs in the theory of Macdonald polynomials. *Proc. Natl. Acad. Sci. USA*, 102(11):3891–3894 (electronic), 2005.

[Jac71] Henry Jack. A class of symmetric polynomials with a parameter. *Proc. Roy. Soc. Edinburgh Sect. A*, 69:1–18, 1970/1971.

[Ker00] S. Kerov. Talk in Institute Henri Poincaré, Paris, January 2000.

[KO94] Serguei Kerov and Grigori Olshanski. Polynomial functions on the set of Young diagrams. *C. R. Acad. Sci. Paris Sér. I Math.*, 319(2):121–126, 1994.

[Las08] Michel Lassalle. A positivity conjecture for Jack polynomials. *Math. Res. Lett.*, 15(4):661–681, 2008.

[Las09] Michel Lassalle. Jack polynomials and free cumulants. *Adv. Math.*, 222(6):2227–2269, 2009.

[LC09] Michael Andrew La Croix. *The combinatorics of the Jack parameter and the genus series for topological maps*. PhD thesis, University of Waterloo, 2009.

[LV95] Luc Lapointe and Luc Vinet. A Rodrigues formula for the Jack polynomials and the Macdonald-Stanley conjecture. *Internat. Math. Res. Notices*, (9):419–424, 1995.

[LV97] Luc Lapointe and Luc Vinet. Rodrigues formulas for the Macdonald polynomials. *Adv. Math.*, 130(2):261–279, 1997.

[Mac95] I. G. Macdonald. *Symmetric functions and Hall polynomials*. Oxford Mathematical Monographs. The Clarendon Press Oxford University Press, New York, second edition, 1995. With contributions by A. Zelevinsky, Oxford Science Publications.

[Śea13] Piotr Śniady and et al. Jack polynomials and the measure of non-orientability of maps. Part 3. Polynomial functions on the set of Young diagrams. In preparation, 2013.

[Sta04] Richard P. Stanley. Irreducible symmetric group characters of rectangular shape. *Sém. Lothar. Combin.*, 50:Art. B50d, 11 pp. (electronic), 2003/04.

[Sta89] Richard P. Stanley. Some combinatorial properties of Jack symmetric functions. *Adv. Math.*, 77(1):76–115, 1989.

[Sta06] Richard P. Stanley. A conjectured combinatorial interpretation of the normalized irreducible character values of the symmetric group. Preprint arXiv:math.CO/0606467, 2006.

INSTYTUT MATEMATYCZNY, UNIWERSYTET WROCŁAWSKI, PL. GRUNWALDZKI 2/4, 50-384 WROCŁAW, POLAND

E-mail address: `Maciej.Dolega@math.uni.wroc.pl`

LABRI, UNIVERSITÉ BORDEAUX 1, 351 COURS DE LA LIBÉRATION, 33 400 TALENCE, FRANCE

E-mail address: `feray@labri.fr`

ZENTRUM MATHEMATIK, M5, TECHNISCHE UNIVERSITÄT MÜNCHEN, BOLTZMANNSTRASSE 3, 85748 GARCHING, GERMANY

INSTYTUT MATEMATYCZNY, POLSKA AKADEMIA NAUK, UL. ŚNIADECKICH 8, 00-956 WARSZAWA, POLAND

INSTYTUT MATEMATYCZNY, UNIWERSYTET WROCŁAWSKI, PL. GRUNWALDZKI 2/4, 50-384 WROCŁAW, POLAND

E-mail address: `piotr.sniady@tum.de, piotr.sniady@math.uni.wroc.pl`